

prolonged disease stabilisation in pre-treated patients who were progressing on the background ET at study entry. Further development of SFX-01 in ER+ mBC is warranted.

**Clinical trial identification:** NCT02970682.

**Legal entity responsible for the study:** Evgen Pharma PLC.

**Funding:** Evgen Pharma PLC.

**Disclosure:** S.J. Howell: Research grant / Funding (self), Travel / Accommodation / Expenses: Evgen Pharma PLC. S. Ross: Full / Part-time employment: Evgen Pharma PLC. T. Morris: Full / Part-time employment: Evgen Pharma PLC. S. Franklin: Leadership role, Shareholder / Stockholder / Stock options, Full / Part-time employment, Officer / Board of Directors: Evgen Pharma PLC. All other authors have declared no conflicts of interest.

**341P Final results of the STEM trial: SFX-01 in the treatment and evaluation of ER+ Her2- metastatic breast cancer (mBC)**

S.J. Howell<sup>1</sup>, M. Campone<sup>2</sup>, J. Cortés<sup>3</sup>, F.P. Duhoux<sup>4</sup>, S. Ross<sup>5</sup>, T. Morris<sup>5</sup>, S. Franklin<sup>5</sup>

<sup>1</sup>Division of Cancer Sciences, The University of Manchester, Manchester, UK, <sup>2</sup>Medical Oncology, ICO Institut de Cancerologie de l'Ouest René Gauducheau, Saint-Herblain, France, <sup>3</sup>Oncology Department, IOB Institute of Oncology, Quironsalud Group, Madrid & Barcelona & Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Madrid, Spain, <sup>4</sup>Medical Oncology, Cliniques Universitaires Saint-Luc, UCLouvain, Brussels, Belgium, <sup>5</sup>Clinical Development, Evgen Pharma PLC, Liverpool, UK

**Background:** SFX-01 is a proprietary synthetic pharmaceutical product based upon a stabilised sulforaphane. In preclinical models SFX-01 inhibits the activity of cancer stem-like cells and reverses resistance to endocrine therapies (ET) tamoxifen (Tam) and fulvestrant (Fulv). The STEM study investigated the potential of SFX-01 to reverse acquired resistance to Tam, Fulv and third generation aromatase inhibitor (AI) therapy.

**Methods:** STEM is an open label parallel arm exploratory phase II trial. Patients were eligible if they had measurable (RECIST v1.1) estrogen receptor (ER) positive Her2-mBC that had developed acquired resistance to AI (arm A), Tam (arm B) or Fulv (arm C) therapy i.e. mBC was progressing having shown clinical benefit (stable disease (SD) for 6 months or objective response (OR)). SFX-01 300mg bd po was added to the current ET. The co-primary endpoints were clinical benefit rate (CBR) and safety/tolerability. Secondary endpoints included OR rate (ORR) and time to progression (TTP). Patients progression free at 24wks were enrolled into a compassionate use phase to continue SFX-01 and the same ET.

**Results:** Between January 2017 and July 2018 46 patients were recruited; arm A n = 31, arm B n = 8 and arm C n = 7. 33/46 (71.7%) had visceral involvement. Prior lines of ET were: 1 = 17/46 (37.0%), 2 = 17/46 (37.0%) and ≥ 3 = 12 (26%). 8 patients (17.4%) had received prior chemotherapy for mBC. The CBR overall was 12/46 (26.0%) with 2 objective responses (both partial). SFX-01 was well tolerated with grade 1/2 nausea (54.3%) and dyspepsia (32.6%) the most frequent adverse events (AEs). Only 2 grade 3 AEs were possibly drug related. There were no drug related serious AEs. There was no significant association between duration of prior ET and subsequent duration of ET+SFX-01 (Spearman  $r$  0.17;  $p$  = 0.26).

**Conclusions:** SFX-01 300 mg BID was safe and well tolerated in patients with ER+ and HER2- mBC. SFX-01 in combination with ET demonstrated anti-tumour activity and